



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Activity Behaviours in Lean and Morbidly Obese Pregnant Women

Citation for published version:

Fazzi gómez, C, Mohd-Shukri, N, Denison, F, Saunders, D, Norman, J & Reynolds, R 2018, 'Activity Behaviours in Lean and Morbidly Obese Pregnant Women', *Scandinavian Journal of Medicine and Science in Sports*. <https://doi.org/10.1111/sms.13219>

Digital Object Identifier (DOI):

[10.1111/sms.13219](https://doi.org/10.1111/sms.13219)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

Scandinavian Journal of Medicine and Science in Sports

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



Title: Activity Behaviours in Lean and Morbidly Obese Pregnant Women.

Authors: Caterina Fazzi¹, Nor Mohd-Shukri², Fiona C Denison¹, David H Saunders³, Jane E Norman¹, Rebecca M Reynolds^{1, 4}.

¹ Tommy's Centre for Maternal and Fetal Health, MRC/University of Edinburgh, Centre for Reproductive Health, Queen's Medical Research Institute, 47 Little France Crescent, Edinburgh EH16 4TJ, United Kingdom.

² Department of Nutrition Sciences, Kulliyyah of Allied Health Sciences, International Islamic University Malaysia, Malaysia.

³ Physical Activity for Health Research Centre (PAHRC), Institute for Sport, Physical Education and Health Sciences, University of Edinburgh.

⁴ University BHF Centre for Cardiovascular Science, University of Edinburgh.

Correspondent Author: Professor Rebecca M Reynolds, Professor of Metabolic Medicine, Centre for Cardiovascular Science, Queen's Medical Research Institute, 47 Little France Crescent, Edinburgh, EH16 4TJ.

Tel: + 44 (0) 131 2426762

Fax: + 44 (01) 131 2426779

e-mail: r.reynolds@ed.ac.uk

Abstract

Interventions to increase physical activity in pregnancy are challenging for morbidly obese women. Targeting sedentary behaviours may be a suitable alternative to increase energy expenditure. We aimed to determine total energy expenditure, and energy expended in sedentary activities in morbidly obese and lean pregnant women.

We administered the Pregnancy Physical Activity Questionnaire PPAQ (non-objective) and the Actical accelerometer (objective) to morbidly obese ($\text{BMI} \geq 40 \text{ kg/m}^2$) and lean ($\text{BMI} \leq 25 \text{ kg/m}^2$) pregnant women recruited in early (< 24 weeks), and late (≥ 24 weeks) gestation. Data are mean (SD).

Morbidly obese pregnant women reported expending significantly more energy per day in early ($n=140$ vs 109 ; 3198.4 (1847.1) vs 1972.3 (10284.8) Kcal/day, $p < 0.0001$) and late ($n=104$ vs 64 ; 3078.2 (1356.5) vs 1947.5 (652.0) Kcal/day, $p < 0.0001$) pregnancy, and expended significantly more energy in sedentary activities, in early (816.1 (423.5) vs 540.1 (244.9) Kcal/day, $p < 0.0001$) and late (881.6 (455.4) vs 581.1 (248.5) Kcal/day, $p < 0.0001$) pregnancy, than lean pregnant women. No differences were observed in the proportion of energy expended sedentary between lean and morbidly obese pregnant women.

The greater total energy expenditure in morbidly obese pregnant women was corroborated by Actical accelerometer in early ($n=14$ per group, obese 1167.7 (313.6) Kcal; lean 781.1 (210.1) Kcal, $p < 0.05$), and in late ($n=14$ per group, obese 1223.6 (351.5) Kcal; lean 893.7 (175.9) Kcal, $p < 0.05$) pregnancy.

In conclusion, non-objective and objective measures showed morbidly obese pregnant women expended more energy per day than lean pregnant. Further studies are needed to determine whether sedentary behaviours are a suitable target for intervention in morbidly obese pregnancy.

Key words: energy expenditure, lean, morbidly obese, pregnancy.

Background

Among women of reproductive age, obesity (body mass index BMI $\geq 30\text{kg/m}^2$) levels have increased in the last decades (1-4). Recent estimates indicate 22% of pregnant women are obese (5), whilst around 2% are severely obese (BMI $\geq 40\text{kg/m}^2$) (6).

Obesity in pregnancy is associated with adverse outcomes for mother and offspring (7, 8). Interventions to increase energy expenditure are an option to control weight and gestational weight gain, but these are challenging to implement in morbidly obese pregnant women (9). Indeed previous studies have shown that levels of physical activity are very low among pregnant women (10), particularly amongst those that are overweight/obese compared with normal-weight (11). Overweight individuals expend considerably more calories than normal weight individuals doing the same exercise (12). Obese pregnant women need more energy to move and have a higher metabolic cost than lean pregnant women, so the work of breathing and moving takes a greater effort, and peripheral motor efficiency is decreased (13). Studies comparing physical activity between obese and normal weight pregnant women are very scarce (11), and the majority of interventions based on increasing physical activity levels in obese pregnant women have been largely unsuccessful in preventing adverse pregnancy outcomes (14-16). Targeting a reduction in sedentary behaviours (i.e., activities that expend very low energy, such as sitting or lying or reclining), may be a realistic alternative (17).

Epidemiological studies show that in the general adult population around 55% to 60% of time awake, is spent sedentary (18, 19). In a systematic review, we showed that pregnant women spend more than 50% of their time sedentary (20). A handful of studies suggest increased time in sedentary behaviours during pregnancy is associated with adverse maternal and offspring outcomes. These include higher maternal levels of LDL cholesterol (21), C-reactive protein (21) and gestational diabetes (22), for the

mother, and higher new born abdominal circumference (23), and risk of macrosomia (birthweight>4000g) (24), for the offspring.

As little is known about sedentary behaviours in morbidly obese pregnant women, we aimed to determine total energy expenditure, and energy expended in sedentary activities in morbidly obese and lean pregnant women using two validated methods, objective (Actical accelerometer) and non-objective (PPAQ). We hypothesised that morbidly obese pregnant women would expend less energy in total activities, but proportionally more time in sedentary activities than lean pregnant women.

Methods

Subjects were morbidly obese ($\text{BMI} \geq 40 \text{ kg/m}^2$) women with a singleton pregnancy attending the Antenatal Metabolic Clinic at the Royal Infirmary of Edinburgh, UK, and lean ($\text{BMI} \leq 25 \text{ kg/m}^2$) pregnant women recruited from community antenatal clinics who were participating in a larger study examining the consequences of morbidly obese pregnancy. Details of the overall cohort have been previously described (25, 26).

Ethical approval was obtained from the Lothian NHS Research Ethics Committee, and all subjects gave informed written consent (REC reference number 08/S1101/39).

In this cross-sectional study women were asked to complete the Pregnancy Physical Activity Questionnaire (PPAQ) in early (<24 weeks' gestation), and late (>24 weeks' gestation) pregnancy. The PPAQ is designed specifically for pregnant women to assess the energy expended in activities of different intensities. It contains 36 questions and was validated against the Actigraph accelerometer (Manufacturing Technology, Inc.) in pregnant women in 2004 (27). Results on energy expenditure are given in metabolic equivalents (28) per day and as total activity plus four different activity levels (sedentary, light, moderate and vigorous). Additionally, energy expenditure is given separately in three type of activities (house activities including caring, occupational, and sports or exercise). To show the data in Kcal per day, we calculated the resting

metabolic rate (RMR) using the Mifflin and St. Jeor equation (29), which has been tested as the best equation to estimate resting energy expenditure in obese and non-obese adults (30, 31).

Energy expenditure was also assessed in early and late pregnancy, in a subset of women (n=14 per group) using the *Actical* accelerometer (Mini Mitter Company, Inc., US), which gives data on Active Energy Expenditure in kilocalories per minute a day, and has been validated for use in healthy adult populations (32). Sedentary activity was classified as time spent performing activities that register less than 100 counts per minute (33). Women wore the device on their non-dominant wrist, for two weekdays and one weekend day, for 24 hours each day (including sleeping time), and were told to remove the Actical only for bathing, or during water sports activities.

Statistical analyses

Data distribution was tested using the Shapiro-Wilk normality test. Continuous variables including time spent in sedentary behaviours, and relative total daily energy expenditure, were compared between morbidly obese and lean pregnant women using T-tests or ANOVA for normally distributed variables and Mann-Whitney U test for data that were not normally distributed. We compared the proportions of energy expended in the different daily activities between groups using ANOVA or Mann-Whitney as appropriate. Regression analyses were used to adjust for potential confounders when analysing the PPAQ. In particular we adjusted for parity and socio-economic status as these have been reported to influence activity levels in other studies (34) and also differed in our sample (supplementary tables 2 and 3). Differences were accepted as significant at $p < 0.05$. Data were analysed using IBM SPSS Statistics 19.0 software.

Results

The PPAQ was completed by 109 lean and 140 morbidly obese women in early pregnancy (<24 weeks, range 12-23 weeks), and 64 lean and 104 morbidly obese women in late pregnancy (\geq 24 weeks, range 24-36 weeks).

Table 1 shows the characteristics of participants who completed the PPAQ. Morbidly obese pregnant women had higher BMI, parity, were of lower deprivation category status, were younger, delivered earlier, and gained significantly less weight than lean pregnant women.

Demographics of the women (n=14 lean early; 14 lean late; n=14 morbidly obese early; 14 morbidly obese late) who wore the accelerometer were similar to the full cohort (Supplementary Table 1).

Total Energy Expenditure and Sedentary Energy Expenditure in morbidly obese and lean pregnant women

When comparing reported energy expenditure using the PPAQ between morbidly obese and lean pregnant women, morbidly obese expended significantly more energy per day as total expenditure and in sedentary activities in both early and late pregnancy, as shown in Table 2. These differences remained significant in regression analyses adjusting for maternal age, parity, deprivation status and ethnicity.

Objective measurements of energy expenditure using the Actical confirmed that morbidly obese pregnant women expended significantly more energy than lean pregnant women in early and late pregnancy despite the observation that in both stages of pregnancy morbidly obese pregnant women performed significantly fewer activity counts than lean pregnant women (Table 3).

Proportions of Total Energy Expenditure in different intensity activities

Proportions of energy expended in different intensities of activity are shown in Figure 1 (a-b). In early and late pregnancy, morbidly obese pregnant women expended

significantly more energy in light intensity and significantly less energy in vigorous intensity activities than lean pregnant women. Differences in the proportion of time spent in vigorous activities remained significant after the regression analysis, controlling for maternal age, parity, deprivation status, and ethnicity. Differences in the proportion of time in light intensity activities did not remain significant in adjusted analyses. No differences were observed between lean and morbidly obese pregnant women in the proportion of time spent in moderate or sedentary intensity activities.

Discussion

Our findings demonstrate that morbidly obese pregnant women expend more energy in all physical activities other than vigorous activities than lean pregnant women. This is despite the observation that morbidly obese pregnant women have fewer objectively measured activity 'counts' than lean pregnant women. Further, though both groups spent a similar time in sedentary activities, morbidly obese pregnant women expended more energy when sedentary than lean pregnant women.

Our observation that morbidly obese pregnant women expended significantly less energy in vigorous activities than lean pregnant women corresponds to other studies showing that this domain of physical activity volume is lower among pregnant women (10), but even lower among overweight or obese pregnant women (11). However, we had anticipated that morbidly obese women would spend proportionally more time in sedentary activities than lean women, but objective measures showed time spent sedentary was similar in both groups. The obese group also expended significantly more total energy daily than lean pregnant women in sedentary activities, consistent with their greater basal metabolic rate (30). Though morbidly obese pregnant women expended significantly more total energy than lean pregnant women, they registered significantly fewer activity counts than lean women using the Actical accelerometer. Counts assessed by Actical are an indication of movement in relation to different planes, gravitational forces, magnitude and duration of the sensed acceleration, but

not linked to personal characteristics such as gender, age, or body weight (33). Thus interventions designed to increase overall movement, many of which could be performed whilst sedentary i.e. sitting, lying, or reclining, may still be a suitable target for morbidly obese pregnant women. Our observations were similar in early and late pregnancy suggesting any intervention should be started in early pregnancy.

A strength of the study is that we used two different methods to assess energy expenditure and sedentary behaviours, including the PPAQ questionnaire, which has been validated in pregnancy, and an objective device. Due to the detailed characterisation of the women we were able to adjust for potential confounding factors including parity and socioeconomic status which were associated with differences in energy expenditure in our sample, as has been reported by others (34). Findings remained significant after adjustment for these confounders. Limitations include the risk of recall bias and potential for lack of reliability of the PPAQ, since subjects might be dishonest or inaccurate in their responses. We also acknowledge the small sample size used with the Actical accelerometer limits interpretation of results. Whilst subjects wore the accelerometer for the recommended time of the manufacturer, we acknowledge this was for a relatively short time. Despite this, the Actical findings for energy expenditure were consistent with the PPAQ outcomes. A further strength is the focus on morbidly obese pregnant women, who may be unable to participate in interventions designed for less severely obese women (15, 35), and have also been identified to have specific barriers to participation in physical activity interventions (36). We acknowledge that time spent sleeping, which may impact on the time spent sedentary, was not specifically assessed in our study, but we are not aware that sleep duration differs between morbidly obese and lean pregnant women (37).

Though we used two validated measures to assess physical activity in pregnancy, neither was specifically designed to understand sedentary activities in pregnancy. A

recent systematic review highlighted the heterogeneity in assessment of sedentary activity (20) with measures ranging from 7 to 18 hours per day.

Perspective

A better understanding of sedentary activity is needed for the design of effective interventions to help to reduce the adverse effects of obesity on pregnancy, especially as obesity prevalence is growing among fertile women (38), and that there are risks associated with obesity during pregnancy, for mothers and offspring. We have shown that morbidly obese pregnant women expend significantly more energy than lean pregnant women, but they also expend significantly more energy on sedentary activities. These findings suggest that energy expenditure might not be the key factor to obesity, but energy intake might be. Nevertheless, sports and physical activity interventions may play a role as preventive health factors contributing to better and effective alternatives to reduce those risks associated with obesity during pregnancy, and to reduce time spent sedentary.

References

1. Heslehurst N, Simpson H, Ells LJ, Rankin J, Wilkinson J, Lang R, et al. The impact of maternal BMI status on pregnancy outcomes with immediate short-term obstetric resource implications: a meta-analysis. *Obes Rev.* 2008;9(6):635-83.
2. Denison FC, Norwood P, Bhattacharya S, Duffy A, Mahmood T, Morris C, et al. Association between maternal body mass index during pregnancy, short-term morbidity, and increased health service costs: a population-based study. *BJOG.* 2014;121(1):72-81; discussion 2.
3. Huda SS, Brodie LE, Sattar N. Obesity in pregnancy: prevalence and metabolic consequences. *Semin Fetal Neonatal Med.* 2010;15(2):70-6.
4. Fitzsimons KJ, Modder J. Setting maternity care standards for women with obesity in pregnancy. *Semin Fetal Neonatal Med.* 2010;15(2):100-7.
5. Scotland ANSPf. Births in Scottish Hospitals. 2016.
6. Enquiries CfMaC. Maternal obesity in the UK: findings from a national project. Executive Summary and Key Recommendations. In: Enquiries CfMaC, editor. United Kingdom 2010.
7. Norman JE, Reynolds RM. The consequences of obesity and excess weight gain in pregnancy. *The Proceedings of the Nutrition Society.* 2011;70(4):450-6.
8. Reynolds RM, Allan KM, Raja EA, Bhattacharya S, McNeill G, Hannaford PC, et al. Maternal obesity during pregnancy and premature mortality from cardiovascular event in adult offspring: follow-up of 1 323 275 person years *BMJ* 2013;347 f4539.
9. Denison FC, Weir Z, Carver H, Norman JE, Reynolds RM. Physical activity in pregnant women with Class III obesity: A qualitative exploration of attitudes and behaviours. *Midwifery.* 2015;31(12):1163-7.
10. Nascimento SL, Surita FG, Godoy AC, Kasawara KT, Morais SS. Physical activity patterns and factors related to exercise during pregnancy: A cross sectional study. *PLoS ONE.* 2015;10(6).

11. Bacchi E, Bonin C, Zanolin ME, Zambotti F, Livornese D, Dona S, et al. Physical Activity Patterns in Normal-Weight and Overweight/Obese Pregnant Women. *Plos One*. 2016;11(11):e0166254.
12. McArdle W, Katch F, Katch V. *Exercise Physiology. Nutrition, Energy, and Human Performance*. Seventh Edition ed. Baltimore: Lippincott Williams & Wilkins; 2010.
13. Mottola MF. Physical activity and maternal obesity: cardiovascular adaptations, exercise recommendations, and pregnancy outcomes. *Nutrition reviews*. 2013;71 Suppl 1:S31-6.
14. Thangaratinam S, Rogozinska E, Jolly K, Glinkowski S, Roseboom T, Tomlinson JW, et al. Effects of interventions in pregnancy on maternal weight and obstetric outcomes: meta-analysis of randomised evidence. *British Medical Journal*. 2012;344.
15. Poston L, Bell R, Croker H, Flynn A, Godfrey K, Goff L, et al. Effect of a behavioural intervention in obese pregnant women (the UPBEAT study): a multicentre, randomised controlled trial. *Lancet Diabetes Endocrinol*. 2015;3(10):767-77.
16. Dodd J, Turnbull D, McPhee A, Deussen A, Grivell R, Yelland L, et al. Antenatal lifestyle advice for women who are overweight or obese: LIMIT randomised trial. *BMJ*. 2014;348.
17. Owen N, Healy GN, Matthews CE, Dunstan DW. Too much sitting: the population health science of sedentary behavior. *Exerc Sport Sci Rev*. 2010;38(3):105-13.
18. Matthews CE, Chen KY, Freedson PS, Buchowski MS, Beech BM, Pate RR, et al. Amount of time spent in sedentary behaviors in the United States, 2003-2004. *Am J Epidemiol*. 2008;167(7):875-81.
19. Spittaels H, Van Cauwenberghe E, Verbestel V, De Meester F, Van Dyck D, Verloigne M, et al. Objectively measured sedentary time and physical activity time across the lifespan. A cross-sectional study in four age groups. *International Journal of Behavioral Nutrition and Physical Activity*. 2012;9(149):1-12.

20. Fazzi C, Saunders DH, Linton K, Norman JE, Reynolds RM. Sedentary Behaviours during Pregnancy: A systematic Review. *International Journal of Behavioral Nutrition & Physical Activity*. 2017;14:32.
21. Loprinzi PD, Fitzgerald EM, Woekel E, Cardinal BJ. Association of physical activity and sedentary behavior with biological markers among U.S. pregnant women. *Journal of Women's Health*. 2013;22(11):953-8.
22. Leng J, Liu G, Zhang C, Xin S, Chen F, Li B, et al. Physical activity, sedentary behaviors and risk of gestational diabetes mellitus: a population-based cross-sectional study in Tianjin, China. *European journal of endocrinology*. 2016;174(6):763-73.
23. Hayes L, Bell R, Robson S, L P. Association between Physical Activity in Obese Pregnant Women and Pregnancy Outcomes: The UPBEAT Pilot Study. *Ann Nutr Metab*. 2014;64:239-46.
24. Reid EW, McNeill JA, Alderdice FA, Tully MA, Holmes VA. Physical activity, sedentary behaviour and fetal macrosomia in uncomplicated pregnancies: A prospective cohort study. *Midwifery*. 2014;30(12):1202-9.
25. Mina TH, Denison FC, Forbes S, Stirrat LI, Norman JE, Reynolds R. Associations of mood symptoms with ante- and postnatal weight change in obese pregnancy are not mediated by cortisol. *Psychol Med*. 2015;45(15):3133-46.
26. Forbes S, Barr SM, Reynolds RM, Semple S, Gray C, Andrew R, et al. Convergence in insulin resistance between very severely obese and lean women at the end of pregnancy. *Diabetologia*. 2015;58(11):2615-26.
27. Chasan-Taber L, Schmidt M, Roberts D, Hosmer D, Markenson G, Freedson P. Development and Validation of a Pregnancy Physical Activity Questionnaire *Medicine and Science in Sports and Exercise*. 2004;36(10):1750-60.
28. Tremblay MS, Warburton DER, Janssen I, Paterson DH, Latimer AE, Rhodes RE, et al. New Canadian Physical Activity Guidelines. *Applied Physiology Nutrition and Metabolism-Physiologie Appliquee Nutrition Et Metabolisme*. 2011;36(1):36-46.

29. Mifflin MD, St Jeor ST, Hill LA, Scott BJ, Daugherty SA, Koh YO. A new predictive equation for resting energy expenditure in healthy individuals. *The American journal of clinical nutrition*. 1990;51(2):241.
30. Frankenfield DC, Rowe WA, Smith JS, Cooney RN. Validation of several established equations for resting metabolic rate in obese and nonobese people. *Journal of the American Dietetic Association*. 2003;103(9):1152-9.
31. Frankenfield D, Roth-Yousey L, Compher C. Comparison of Predictive Equations for Resting Metabolic Rate in Healthy Nonobese and Obese Adults: A Systematic Review. *Journal of the American Dietetic Association*. 2005;105(5):775-89.
32. Crouter SE, Dellavalle DM, Horton M, Haas JD, Frongillo EA, Bassett DR. Validity of the Actical for estimating free-living physical activity. *Eur J Appl Physiol*. 2010;111(7):1381-9.
33. Lai S, Colley R, Connor S, M T. Actical Accelerometer Sedentary Activity Thresholds for Adults. *Journal of Physical Activity and Health*. 2011;8:587 -91.
34. Gaston A, Cramp A. Exercise during pregnancy: a review of patterns and determinants. *J Sci Med Sport*. 2011;14(4):299-305.
35. Dodd JM. Dietary and lifestyle advice for pregnant women who are overweight or obese: the LIMIT randomized trial. *Annals of nutrition & metabolism*. 2014;64(3-4):197-202.
36. Weir Z, Bush J, Robson SC, McParlin C, Rankin J, R B. Physical activity in pregnancy: a qualitative study of the beliefs of overweight and obese pregnant women. *BMC Pregnancy and Childbirth*. 2010;10.
37. Qiu C, Frederick IO, Sorensen TK, Enquobahrie DA, MA W. Sleep duration and plasma leptin concentrations in early pregnancy among lean and overweight/obese women: a cross sectional study. *BMC Research Notes*. 2014;7:20.
38. Fisher SC, Kim SY, Sharma AJ, Rochat R, B. M. Is obesity still increasing among pregnant women? Prepregnancy obesity trends in 20 states, 2003–2009. *Prev Med*. 2013;56(6):372-8.

39. Szymanski LM, Satin AJ. Exercise During Pregnancy Fetal Responses to Current Public Health Guidelines. *Obstet Gynecol.* 2012;119(3):603-10.
40. Owen N, Leslie E, Salmon J, Fotheringham MJ. Environmental Determinants of Physical Activity and Sedentary Behavior. *Exercise and Sport Sciences Reviews.* 2000;28(4):153-8.

Tables.**Table 1. Characteristics of obese and lean participants who completed the PPAQ in early and late pregnancy.**

| Characteristic | EARLY | | | LATE | | |
|--|---|---|----------------|--|---|----------------|
| | Lean (n=109) Mean (SD) or n (%) | M. Obese (n=140) Mean (SD) or n (%) | p-value | Lean (n=64) Mean (SD) or n (%) | M. Obese (n=104) Mean (SD) or n (%) | p-value |
| BMI (Kg/m ²) | 22.8 (2.7) | 44.2 (4.5) | <0.001 | 22.8 (1.6) | 44.1 (5.0) | <0.001 |
| Maternal age (years) | 33.06 (4.55) | 30.73 (5.40) | <0.001 | 33.61 (4.45) | 31.50 (5.26) | <0.05 |
| Parity | | | <0.01 | | | <0.01 |
| 0 | 68 (62.4) | 64 (46) | | 41 (64.1) | 43 (41) | |
| 1 | 29 (26.6) | 41 (29.5) | | 16 (25) | 38 (36.2) | |
| 2 | 12 (11) | 31 (39) | | 7 (10.9) | 20 (19) | |
| 3 | 0 (0) | 2 (1.4) | | 0 (0) | 2 (1.9) | |
| 4 | 0 (0) | 0 (0) | | 0 (0) | 0 (0) | |
| 5 | 0 (0) | 0 (0) | | 0 (0) | 2 (1.9) | |
| Ethnicity | | | >0.05 | | | >0.05 |
| Caucasian | 97 (89) | 124 (89.2) | | 54 (84.4) | 84 (81) | |
| Other | 0 (0) | 4 (2.9) | | 0 (0) | 3 (2.9) | |
| Deprivation Category | | | <0.001 | | | <0.01 |
| Low | 28 (25.9) | 11 (8.0) | | 15 (24.6) | 12 (11.7) | |
| Middle | 79 (73.1) | 103 (75.7) | | 46 (75.4) | 79 (76.7) | |
| High | 1 (0.9) | 22 (16.2) | | 34 (0) | 12 (11.7) | |
| Birth weight (g) | 3513 (541) | 3574 (558) | >0.05 | 3584 (512) | 3511(595) | >0.05 |
| Gestational age at delivery (week) | 40.34 (1.34) | 39.79 (1.50) | <0.005 | 40.50 (1.38) | 39.68 (1.42) | <0.001 |
| Weight gain (kg) | 10.16 (3.64) | 5.87 (5.03) | <0.001 | 10.41 (4.05) | 5.59 (5.53) | <0.001 |

Data are mean (SD) or n (%). Weight gain was calculated as Weight week 36 –weight weight at baseline.
Deprivation Category is based on postcodes.

Table 2. PPAQ comparisons in total and sedentary energy expenditure between lean and morbidly obese pregnant women in early and late stage.

| | Early | | | Late | | |
|--|----------------------|----------------------|----------------------------|---------------------|----------------------|----------------------------|
| | Mean (SD) | | β (95% CI) † | Mean (SD) | | β (95% CI) † |
| | Lean (n=109) | M.Obese (n=140) | | Lean (n=64) | M.Obese (n=104) | |
| Total EE (Kcal/day) | 1972.29 (1028.85) | 3198.37 (1847.05) | 0.33**(575.73 -1390.80) | 1947.54 (652.03) | 3078.23 (1356.46) | 0.43**(699.87 -1631.39) |
| Sedentary Activity EE (Kcal/day) | 590.13 (244.90) | 816.07 (423.51) | 0.37**(180.08 -397.11) | 581.11 (248.51) | 881.65 (455.38) | 0.34**(110.69 -360.39) |

β is the standardised coefficient.

**Significant at $p < 0.001$

†Adjusted for Maternal Age, Parity, Deprivation Category, and Ethnicity.

Table 3. Actical comparisons in counts, total energy expenditure, and sedentary time between lean and morbidly obese pregnant women in early and late stage.

| | Early Mean (SD) | | | Late Mean (SD) | | |
|--|--------------------------|-------------------------|------------|-------------------------|-------------------------|------------|
| | Lean (n=14) | M.Obese (n=14) | Sig | Lean (n=14) | M.Obese (n=14) | |
| Sedentary time (min/day) | 762.40 (104.68) | 799.33 (101.80) | $p > 0.05$ | 740.70 (89.89) | 774.15 (124.70) | $p > 0.05$ |
| Total Activity Counts* (per day) | 360160.91 (131302.13) | 268683.36 (83567.16) | $p < 0.05$ | 357561.03 (94799.09) | 266820.25 (97640.51) | $p < 0.05$ |
| Total EE (Kcal/day) | 781.06 (210.15) | 1167.69 (313.56) | $p < 0.01$ | 893.72 (175.88) | 1223.64 (351.47) | $p < 0.05$ |

*Counts are markers of movement.

Supplementary Tables.

Supplementary Table 1. Characteristics of Actical accelerometer participants by groups.

| Characteristic | Lean Early* (n=14) Mean (SD) | M. Obese Early* (n=14) Mean (SD) | p-value | Lean Late† (n=14) Mean (SD) | M. Obese Late† (n=14) Mean (SD) | p-value |
|--------------------------|------------------------------------|--|---------|-----------------------------------|---------------------------------------|---------|
| Maternal Age (years) | 31.08 (4.96) | 31.43 (5.11) | >0.05 | 34.62 (4.81) | 34.86 (4.19) | >0.05 |
| BMI (Kg/m ²) | 23.44 (1.18) | 43.65 (2.99) | <0.001 | 25.81 (2.13) | 44.00 (2.69) | <0.001 |
| % Fat mass | 30.08 (3.76) | 49.26 (1.58) | <0.001 | 33.09 (3.24) | 50.36 (2.23) | <0.001 |
| Parity | | | >0.05 | | | >0.05 |
| 0 | 9 (64.3) | 6 (42.9) | | 7 (50) | 6 (42.9) | |
| 1 | 4 (28.6) | 4 (28.6) | | 6 (42.9) | 3 (21.4) | |
| 2 | 0 (0) | 3 (21.4) | | 1 (7.0) | 2 (14.3) | |
| 3 | 1 (7.1) | 0 (0) | | 0 (0) | 1 (7.1) | |
| 4 | 0 (0) | 1 (7.1) | | 0 (0) | 1 (7.1) | |
| 5 | 0 (0) | 0 (0) | | 0 (0) | 1 (7.1) | |
| Ethnicity | | | >0.05 | | | >0.05 |
| 1 (Caucasian) | 14 (100) | 12 (85.71) | | 14 (100) | 12 (85.71) | |
| 2 (40) | 0 (0) | 2 (14.29) | | 0 (0) | 2 (14.29) | |
| Deprivation Category | | | <0.05 | | | <0.05 |
| Low | 7 (50) | 1 (7.15) | | 4 (28.6) | 0 (0) | |
| Middle | 7 (50) | 12 (85.7) | | 10 (71.4) | 11 (78.6) | |
| High | 0 (0) | 1 (7.15) | | 0 (0) | 3 (21.4) | |
| Birthweight (gr) | 3844.73 (463.88) | 3581.75 (763.34) | >0.05 | 3910.00 (485.02) | 3819.50 (421.38) | >0.05 |
| Weight Gain (Kg) | 10.39 (4.92) | 5.49 (1.91) | =0.058 | 12.19 (3.82) | 7.44 (6.05) | <0.05 |
| BMR (Kcal/day) | 1442.79 (75.20) | 1894.21 (97.08) | <0.001 | 1496.71 (86.79) | 1929.57 (106.06) | <0.001 |

*Early gestation is between 14 and 23 weeks (median 17.93 weeks).

†Late gestation is between 27 and 37 weeks (median 29.36).

Supplementary Table 2. Comparisons on energy expenditure between nulliparous and multiparous pregnant women.

| | EARLY Mean (SD) | | | LATE Mean (SD) | | |
|------------------------|------------------------|------------------------|-----------------|-----------------------|-----------------------|-----------------|
| | Nulliparous (n=132) | Multiparous (n=117) | <i>p</i> -value | Nulliparous (n=84) | Multiparous (n=84) | <i>p</i> -value |
| Total EE (Kcal/day) | 2178.16 (1133.9) | 3207.14 (1962.1) | <0.001 | 2367.70 (1218.1) | 2927.29 (1254.6) | <0.001 |
| Sed EE (Kcal/day) | 762.14 (335.7) | 666.43 (406.9) | <0.01 | 811.03 (474.6) | 723.28 (343.7) | >0.05 |
| Light EE (Kcal/day) | 688.46 (461.4) | 1274.69 (654.0) | <0.001 | 754.42 (481.8) | 1225.42 (561.6) | <0.001 |
| Mod EE (Kcal/day) | 695.76 (791.4) | 1236.79 (1351.4) | <0.001 | 780.77 (864.2) | 956.04 (810.7) | <0.01 |
| Vig EE (Kcal/day) | 30.58 (59.4) | 29.34 (67.6) | >0.05 | 21.62 (49.0) | 21.94 (54.4) | >0.05 |


Supplementary Table 3. Comparisons on energy expenditure between most deprived and least deprived pregnant women.

| | EARLY Mean (SD) | | | LATE Mean (SD) | | |
|------------------------|--------------------------|--------------------------|-----------------|--------------------------|--------------------------|-----------------|
| | Most Deprived (n=167) | Least Deprived (n=78) | <i>p</i> -value | Most Deprived (n=110) | Least Deprived (n=53) | <i>p</i> -value |
| Total EE (Kcal/day) | 2531.19 (1608.6) | 2844.74 (1598.0) | <0.05 | 2588.14 (1335.9) | 2816.03 (1133.9) | >0.05 |
| Sed EE (Kcal/day) | 733.44 (367.9) | 689.58 (387.1) | >0.05 | 769.92 (452.7) | 756.66 (336.6) | >0.05 |
| Light EE (Kcal/day) | 878.67 (604.6) | 1131.62 (654.5) | <0.01 | 942.24 (572.4) | 1106.75 (569.2) | >0.05 |
| Mod EE (Kcal/day) | 892.70 (1065.1) | 986.13 (1047.2) | >0.05 | 853.58 (925.4) | 932.70 (668.7) | >0.05 |
| Vig EE (Kcal/day) | 25.45 (47.3) | 37.50 (87.2) | >0.05 | 22.24 (51.8) | 19.49 (49.9) | >0.05 |

Figures

Figure 1.


a) Percentage of self-reported Energy Expenditure per Activity Intensity in early pregnancy.



*Significant at $p < 0.05$

**Significant at $p < 0.001$

b) Percentage of self-reported Energy Expenditure per Activity Intensity in late pregnancy.



*Significant at $p < 0.05$

**Significant at $p < 0.001$